

SOIL SAMPLING

A CRITICAL ACTIVITY FOR NUTRIENT MANAGEMENT



INTRODUCTION

Soil sampling assists in determining suitable management strategies to optimise crop performance. Different assays are available to assess a range of important soil chemical, physical and biological properties. This article focuses on soil sampling to determine crop nutritional requirements consistent with the SIX EASY STEPS™ approach.

WHY SOIL SAMPLE?

- Diagnose nutritional problems and soil chemical constraints (such as acidity, sodicity, salinity)
- Determine crop nutrition requirements
- Reduce the risk of under-fertilising and cost of over-fertilising
- Monitor the effectiveness of nutrition and amelioration programs
- Optimise other crop management practices by understanding soil properties

HOW TO SOIL SAMPLE?

There are five important steps in the soil sampling process

Step 1: Planning

Review available information to develop a **'fit for purpose' soil sampling strategy**. Identify which blocks need to be sampled and consider soil variability, position in the landscape, topography, crop performance and management history. Sampling each block individually will provide a more accurate assessment but it may be

possible to rationalise the number of samples due to common soil types and management history.

Following this assessment, the most suitable soil sampling pattern, can be identified for each block. Try to collect soil samples at the same time each year, avoid sampling immediately after high

rainfall or irrigation and delay sampling for at least two months after applying any ameliorants or fertiliser.

Gather the right equipment - soil sampling plan, soil auger or probe, plastic handled bucket, clean plastic bag/s and permanent marker.

Step 2: Collection

Clean hands and equipment. To avoid sample contamination wash hands prior to sampling. Ensure equipment is clean (to manage farm hygiene sanitise with 70% methylated spirits) and hasn't been in contact with fertiliser.


Avoid using or touching galvanised surfaces (like bucket

handles and vehicle trays) during sample collection to prevent contamination.


Collect samples from the correct position on the row – this will be influenced by fertiliser placement in the previous crop. Generally, soil samples are collected from the shoulder of the cane row

(about halfway between the centre of the cane row and centre of the inter-row). If fertiliser was applied using a side-dress applicator, soils cores need to be collected closer to the centre of the sugarcane row rather than the shoulder. For reliable results, avoid collecting samples from non-cropped areas,

Single row



Dual row



poorly drained sections and areas where fertiliser or soil ameliorants including mill by-products have been stored. Also avoid including crop residue or organic matter from the soil surface in the sample.

Collect samples from the correct soil sampling depth. To determine crop nutritional requirements sample the topsoil (0-20cm). Other soil depths may be sampled to support the identification of subsoil constraints.

To ensure a representative sample, collect an appropriate number of soil cores from the sampling zone. This may be a whole block, portion of a block or pre-determined zones within a block. Generally, the more soil cores taken, the more reliable the soil test result. As a rule of thumb collect about 20 soil cores from each sampling zone. This can be varied depending on the area of the sampling zone.

A common mistake is to collect too few cores to adequately represent the zones being tested, particularly in very large blocks. In some cases, it is sensible to divide larger areas into smaller sampling zones. Electromagnetic (EM) mapping can also be used to support the identification of suitable soil sampling zones.


Thoroughly mix soil cores, taking care to break-up large soil clods as you blend the sample by hand.

Collect a subsample (500g) from the blended soil sample for laboratory analysis. More soil will be required for a subsample if requesting additional soil assays. Place the subsample into a clean plastic bag, such as those supplied by soil testing laboratories or a ziplock sandwich bag. Clearly label each subsample

– include date, name, farm number, block number or sampling location (if taking more than one sample per block).

Record soil sampling event details – including dominant soil type/s and geo-location of soil sampling sites.

This will allow variations in soil properties to be better understood and the same sampling locations to be revisited overtime. This is valuable for monitoring changes in soil fertility and checking on the adequacy of nutrient inputs.



6,000 tonnes of soil in a 2 ha sample area




Figure 1: Importance of correct soil sampling procedure

Step 3: Sample handling

Soil samples should be dispatched to the laboratory as soon as possible after collection. For short term storage, soil samples should be placed in an esky or refrigerator. Do not leave samples inside a vehicle in the sun.

Step 4: Laboratory analysis

Samples should be submitted to a reputable laboratory for analysis through agribusiness agents, productivity services organisations (in some districts) or private agronomic consultants. The laboratory must conduct tests and report results that are compatible with the SIX EASY STEPS® guidelines. Ensure the laboratory used has a Certificate of Proficiency issued by the Australasian Soil and Plant Analysis Council (ASPAC).

Refer to the *Analyses required to interpret a soil test using the SIX EASY STEPS* information sheet available on the Nutrient Management webpage of the SRA website. Or scan the QR code on the right.



Step 5: Interpretation of results

Soil test reports should be interpreted using the SIX EASY STEPS nutrient management guidelines relevant to your region. This is best done in conjunction with your trusted advisor.

Information pertaining to fallow management and application of mill by-products is important for interpreting soil test results and providing sound nutrient management advice.

Participating in a SIX EASY STEPS workshop or Online Sugarcane Nutrient Management training program will equip you with the knowledge to develop block specific soil sampling strategies, confidently collect soil samples, interpret soil test reports, identify suitable fertiliser products and develop a basic nutrient management plan for your farm.

If interested in SIX EASY STEPS training contact your SRA District Manager.

Acknowledgement:

This information sheet was developed in collaboration with Professor Bernard Schroeder from the University of Southern Queensland and Sugar Research Australia through funding from the Australian Government's National Landcare Program and the Cassowary Coast Reef Smart Farming project funded by the partnership between the Australian Government's Reef Trust and the Great Barrier Reef Foundation.



Copyright © 2024 • All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without the prior permission of SRA. **Disclaimer:** In this disclaimer a reference to 'we', 'us' or 'our' means SRA and our directors, officers, agents and employees. Although we do our best to present information that is correct and accurate, we make no warranties, guarantees or representations about the suitability, reliability, currency or accuracy of the information we present in this Information Sheet, for any purposes. Subject to any terms implied by law and which cannot be excluded, we accept no responsibility for any loss, damage, cost or expense incurred by you as a result of the use of, or reliance on, any materials and information appearing in this Information Sheet. You, the user, accept sole responsibility and risk associated with the use and results of the information appearing in this Information Sheet, and you agree that we will not be liable for any loss or damage whatsoever (including through negligence) arising out of, or in connection with the use of this Information Sheet. We recommend that you contact our staff before acting on any information provided in this Information Sheet. **Warning:** Our tests, inspections and recommendations should not be relied on without further, independent inquiries. They may not be accurate, complete or applicable for your particular needs for many reasons, including (for example) SRA being unaware of other matters relevant to individual crops, the analysis of unrepresentative samples or the influence of environmental, managerial or other factors on production.